

CP violation in $B_{d,s} \rightarrow l^+ l^-$

Chao-Shang Huang (Presenting author)^a, Wei Liao^b

^aInstitute of Theoretical Physics, Academia Sinica, 100080 Beijing, China

^bThe Abdus Salam International Centre for Theoretical Physics, Trieste, Italy

CP violation in rare leptonic B decays are analyzed in the standard model (SM) and models beyond SM (supersymmetric models and two Higgs doublet models).

1. INTRODUCTION

The FCNC process $B_{d,s}^0 \rightarrow l^+ l^-$ has been shown in recent years to be a powerful process to shed light on new physics beyond SM [1,2,3,4,5] especially for SUSY models which may enhance the decay amplitude by $\tan^3 \beta$ [1,6,2,3], provided that $\tan \beta$ is large (say, ≥ 20). It became more interesting recently after the Brookhaven National Laboratory (BNL) reported the 2σ excess of the muon anomalous magnetic moment $a_\mu = (g - 2)_\mu/2$ over its SM value: $\Delta a_\mu \equiv a_\mu^{exp} - a_\mu^{SM} = 2 \times 10^{-9}$ at a 2σ level [7]. SUSY models with large $\tan \beta$ (say, ≥ 10) are favored by this excess. It was shown for mSUGRA in [4] that with a good fit to $(g - 2)_\mu$ ¹ the branching ratio (Br) of $B_{d,s}^0 \rightarrow l^+ l^-$ can be enhanced by 100 times and within good reach at Tevatron Run II.

Besides Br, is there CP violation in the decays which can be observed? In this talk I shall discuss the subject.

CP violation in the b -system has been established from measurements of time-dependent asymmetries in $B \rightarrow J/\Psi K$ decays [8,9]. We shall show that to observe the CP asymmetry in the B decays to a pair of muons or taus is a good way to search for new physics. Direct CP violation is absent and no T-odd projections can be defined in the decays. However, there is CP violation induced by $B^0 - \bar{B}^0$ mixing in the process

$$B^0 \rightarrow \bar{B}^0 \rightarrow f \quad vs. \quad \bar{B}^0 \rightarrow B^0 \rightarrow \bar{f}.$$

¹The analysis in ref.[4] was based on the old data, $\Delta a_\mu \equiv a_\mu^{exp} - a_\mu^{SM} = (43 \pm 16) \times 10^{-10}$.

One can define the CP violating observable as

$$\begin{aligned} A_{CP} &= \frac{D}{S}, \\ D &= \int_0^\infty dt \sum_{i=1,2} \Gamma(B_{phys}^0(t) \rightarrow f_i) \\ &\quad - \int_0^\infty dt \sum_{i=1,2} \Gamma(\bar{B}_{phys}^0(t) \rightarrow \bar{f}_i), \\ S &= \int_0^\infty dt \sum_{i=1,2} \Gamma(B_{phys}^0(t) \rightarrow f_i) \\ &\quad + \int_0^\infty dt \sum_{i=1,2} \Gamma(\bar{B}_{phys}^0(t) \rightarrow \bar{f}_i) \end{aligned} \quad (1)$$

where $f_{1,2} = l_{L,R}^+ l_{L,R}^-$ with $l_{L(R)}$ being the helicity eigenstate of eigenvalue $-1(+1)$, $\bar{f}_i$ is the CP conjugated state of f_i .

In SM, one has

$$\frac{q}{p} = -\frac{M_{12}^*}{|M_{12}|} = -\frac{\lambda_t^*}{\lambda_t}, \quad (2)$$

where $\lambda_t = V_{tb}V_{td}^*$ or $V_{tb}V_{ts}^*$, up to the correction smaller than or equal to order of 10^{-2} , it is

$$A_{CP} = -\frac{2Im(\xi)X_q}{(1+|\xi|^2)(1+X_q^2)}, \quad q = d, s, \quad (3)$$

where $X_q = \frac{\Delta m_q}{\Gamma}$ ($q = d, s$ for B_d^0 and B_s^0),

$$\xi = \frac{C_{Q1}\sqrt{1-4\hat{m}_l^2} + (C_{Q2} + 2\hat{m}_l C_{10})}{C_{Q1}^*\sqrt{1-4\hat{m}_l^2} - (C_{Q2}^* + 2\hat{m}_l C_{10}^*)}. \quad (4)$$

In eq. (4) $\hat{m}_l = m_l/m_{B^0}$. In SM C_{10} is real, $C_{Q2} = 0$, and C_{Q1} is negligibly small. Therefore, there is no CP violation in SM. If one includes the correction of order of 10^{-2} to $|\frac{q}{p}| = 1$,

one will have CP violation of order of 10^{-3} which is unobservably small. It should be pointed out that there is no hadronic uncertainty since the common uncertain decay constant, which is the only source of hadronic uncertainty for the process, cancels out in eq. (4). Because C_{Q_i} 's are proportional to m_l and C_{10} is independent of m_l it follows from eq. (4) that CP asymmetry in $B_{d,s} \rightarrow l^+ l^-$ is independent of the mass of the lepton, i. e., it is the same for $l = \text{electron, muon and tau}$, when one assumes $\sqrt{1 - 4\hat{m}_l^2} \simeq 1$ which is of a good approximation for electron and muon.

In MSSM we can still use eq. (3) in the approximation, eq. (2), which is a good approximation in MSSM if one limits himself to the regions with large $\tan\beta$ (say, larger than 10 but smaller than 60), not too light charged Higgs boson (say, larger than 250 GeV), and heavy sparticles, and in the scenarios of the minimal flavor violation (MFV)² without new CP violating phases there is no correction to eq. (2) [10,11]. In MFV models with new CP violating phases, e.g., in the CMSSM with nonuniversal gaugino masses which we consider in this talk, a rough estimate gives that the correction to the SM value of q/p is below 20% in the parameter space we used in calculations [16].

From eq. (4), we need to calculate the Wilson coefficients C_{Q_i} ($i=1,2$) and C_{10} in models beyond SM in order to compute the CP asymmetry. We shall consider the CP spontaneously broken 2HDM [12] and the CMSSM with nonuniversal gaugino masses [13]. Compared with mSUGRA with CP violating phases (the phase of the Higgsino mass parameter μ and the phase of A), in the CMSSM with nonuniversal gaugino masses there are two more real parameters (say, $|M_1|$ and $|M_3|$, where M_1 and M_3 are gaugino masses corresponding to $U(1)$ and $SU(3)$ respectively) and two more independent phases arising from complex gaugino masses, which make the cancellations among various SUSY contributions to EDMs easier than in mSUGRA with CP violating phases and relatively large values of the phase of μ are allowed [17]. The Wilson coefficients

C_{Q_i} ($i=1,2$) and C_{10} in 2HDM have been calculated respectively in ref.[14,12] and ref.[15]. In CMSSM they have also been calculated and can be found in refs. [1,2,3]³. We show the leading terms of relevant Wilson coefficients in the large $\tan\beta$ case in CMSSM in the following.

$$C_{10}(m_W) = \frac{m_\ell^2}{m_W^2} \tan^2 \beta \sqrt{x_{\chi_j^-} x_{\chi_i^-}} U_{j2}^* U_{i2} f_{D^0}(x_{\chi_i^-}, x_{\chi_j^-}, x_{\tilde{u}_k}, x_{\tilde{\nu}_l}) + \dots \quad (5)$$

$$C_{Q_1}(m_W) = -\tan^3 \beta \frac{m_b m_\ell}{4 \sin^2 \theta_w m_W \lambda_t} \sum_{i=1}^2 \sum_{k=1}^6 U_{i2} T_{UL}^{km} K_{mb} \{ -\sqrt{2} V_{i1}^* (T_{UL} K)_{ks} + V_{i2}^* \frac{(T_{UR} \tilde{m}_u K)_{ks}}{m_W \sin \beta} \} r_{hH} \sqrt{x_{\chi_i^-}} f_{B^0}(x_{\chi_i^-}, x_{\tilde{u}_k}) + O(\tan^2 \beta), \quad (6)$$

$$C_{Q_2}(m_W) = \tan^3 \beta \frac{m_b m_\ell}{4 \sin^2 \theta_w m_W \lambda_t} \sum_{i=1}^2 \sum_{k=1}^6 U_{i2} T_{UL}^{km} K_{mb} \{ -\sqrt{2} V_{i1}^* (T_{UL} K)_{ks} + V_{i2}^* \frac{(T_{UR} \tilde{m}_u K)_{ks}}{m_W \sin \beta} \} r_A \sqrt{x_{\chi_i^-}} f_{B^0}(x_{\chi_i^-}, x_{\tilde{u}_k}) + O(\tan^2 \beta), \quad (7)$$

where U and V are matrices which diagonalize the mass matrix of charginos, T_{Ui} ($i=L, R$) is the matrix which diagonalizes the mass matrix of the scalar up-type quarks and K is the CKM matrix.

The $\tan^3 \beta$ enhancement of C_{Q_i} ($i=1,2$) was first shown in ref. [1] and confirmed later in refs. [6,2,3]. The chargino-chargino box diagram gives a contribution proportional to $\tan^2 \beta$ to C_{10} . Numerically, C_{10} is enhanced in CMSSM at most by about 10% compared with SM.

³Note that the explicit expressions of the Wilson coefficients C_{Q_i} 's are the same in SUSY models with and without CP violating phases. In the SUSY models with CP violating phases the coefficients become complex since the new CP violating phases enter into squark and chargino mass matrices [16].

²MFV means the models in which the CKM matrix remains the unique source of flavor violation.

In a 2HDM with CP violating phases and CMSSM with nonuniversal gaugino masses the CP asymmetries depend on the parameters of models and can be as large as 40% for B_d^0 and 3% for B_s^0 , while the constraints from EDMs of electron and neutron are satisfied [16].

The correlations between $(g - 2)_\mu$ and CP asymmetries in $B_{d,s}^0 \rightarrow l^+ l^-$ and $b \rightarrow s\gamma$ in SUSY models with nonuniversal gaugino masses have been calculated in ref. [16], imposing the constraints from the branching ratio of $B \rightarrow X_s \gamma$ (it leads to the correlation between C_7^{eff} and C_{Q_i} in SUSY models) and EDMs of electron and neutron, and the results are

—with a good fit to the muon $g - 2$ constraint, the CP asymmetry can be as large as 25% (15%) for $B_d^0 \rightarrow \tau^+ \tau^-$ ($B_d^0 \rightarrow \mu^+ \mu^-$) in CMSSM with nonuniversal gaugino masses and MFV scenarios of MSSM.

—If tau events identified with 6% tagging error, one can measure A_{CP} to a 3σ level at Tevatron Run II with $\text{Br}(B_d \rightarrow \tau^+ \tau^-)$ enhanced by a factor of about 30 compared to that of SM.

—A scenario in which new physics only increases the Br a little and the Br is still in the uncertain region of the SM prediction. The CP asymmetry for $B_d^0 \rightarrow l^+ l^-$ in the scenario can still reach 20% allowed by the muon $g-2$ constraint within 2σ deviations. So it is powerful to shed light on physics beyond SM while the CP asymmetry of $b \rightarrow s\gamma$ in this case can only reach 2% at most which is too small to draw a definite conclusion on new physics effects at B factories.

In summary, an observation of CP asymmetry in the decays $B_q^0 \rightarrow l^+ l^-$ ($q = d, s, l = \mu, \tau$) would unambiguously signal the existence of new physics.

REFERENCES

1. C.-S. Huang and Q.-S. Yan, Phys. Lett. **B442** (1998) 209; C.-S. Huang, W. Liao and Q.-S. Yan, Phys. Rev. **D59** (1999) 011701.
2. K.S. Babu and C. Kolda, Phys. Rev. Lett. **84** (2000) 228.
3. C.-S. Huang et al., Phys. Rev. **D63** (2001) 114021; *ibid.* **64** (2001) (2001) 059902(E); P.H. Chankowski, L. Slawianowska, P.R. **D63** (2001) 054012; C. Bobeth et al., Phys. Rev. **D64** (2001) 074014; hep-ph/0204225; G. Isidori, A. Retico, JHEP **11** (2001) 001.
4. A. Dedes, H. K. Dreiner and U. Nierste, Phys. Rev. Lett. **87** (2001) 251804.
5. R. Arnowitt et al., hep-ph/0203069; S. Baek, P. Ko and W. Y. Song, hep-ph/0205259; H. Dreiner, U. Nierste and P. Richardson, hep-ph/0207026; J. K. Mizukoshi, X. Tata and Y. Wang, hep-ph/0208078. For a recent review, see, e.g., Chao-Shang Huang, hep-ph/0210314.
6. C. Hamzaoui, M. Pospelov and M. Toharia, Phys. Rev. **D59** (1999) 095005.
7. G.W. Bennet *et al.*, hep-ex/0208001.
8. B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **87**, 091801 (2001).
9. K. Abe *et al.* [Belle Collaboration], Phys. Rev. Lett. **87**, 091802 (2001).
10. A.J. Buras et al., hep-ph/0107048.
11. G. Isidori and A. Retico, hep-ph/0110121.
12. C.-S. Huang and S.-H. Zhu, Phys. Rev. **D61** (2000) 015011, Erratum-*ibid.* **D61** (2000) 119903. Chao-Shang Huang, Liao Wei, Qi-Shu Yan, Shou-Hua Zhu, Euro. Phys. J. **C25** (2002) 103.
13. A. Brignole, L. Ibáñez, C. Muñoz, Nucl. Phys. **B422** (1994) 125, *ibid.* **B436** (1995) 747(E); M.Brhlík, G.J.Good, and G.L.Kane, Phys. Rev. **D59** 11504(1999); N. Chamoun, C.-S. Huang, C. Liu and X.-H. Wu, Nucl. Phys. **B624** (2002) 81.
14. Y.-B. Dai, C.-S. Huang, and H.-W. Huang, Phys. Lett. **B390** (1997) 257; *ibid.* **B513** (2001) 429(E); H.E. Logan and U. Nierste, Nucl. Phys. **B586** (2000) 39.
15. B. Grinstein, M.J. Savage, M.B. Wise, Nucl. Phys. **B319** (1989) 271.
16. C.-S. Huang, W. Liao, Phys. Lett. **B525** (2002) 107; Phys. Lett. **B538** (2002) 301.
17. T. Ibrahim, P. Nath, Phys. Lett **B418** (1998) 98; C.-S. Huang, W. Liao, Phys. Rev. **D62** (2000) 016008.